

# The Gastrointestinal System and Oral PK in Göttingen Minipigs

The pig is an omnivore and has a digestive physiology very similar to that in humans. Accordingly, Göttingen Minipigs have been used in nutritional studies; in some cases, fitted with re-entrant fistulas to study regional intestinal digestion.<sup>1</sup> The comparative biology of the gastrointestinal system in Minipigs and humans, including specific anatomic differences, have been thoroughly described elsewhere.<sup>2-5</sup> The characteristics of the different segments of the gastrointestinal tracts (i.e. size, pH, etc.) are nicely summarized by Suenderhauf et al.<sup>4</sup>


As a reflection of the clinical situation, the oral route of administration is the most commonly used in toxicology studies (see the illustration below). There are several ways of dosing orally in Göttingen Minipigs but in toxicology studies, the compound is usually given by gastric lavage using a volume of 2-10 ml/kg - in accordance with the administration volumes considered good practice.<sup>6</sup> A dosing chair, as the one described in Newsletter No. 47 (for download at [www.minipigs.dk](http://www.minipigs.dk)), can be of help and depending on the technique used and the size of the Minipigs, two to three persons are usually involved in dosing a group of Minipigs.

The gastric pH in the fasted Minipig is less variable and more comparable to that in humans than in the fasted dog (where it is

often basic).<sup>4,5,7</sup> The gastric emptying of the Minipig is, however, slower than the gastric emptying observed in humans.<sup>8,9</sup> By removing the straw bedding (and potentially replacing it with other enrichment types), the stomach content remaining after an overnight fast can be markedly reduced.<sup>8</sup> In such a setting, mean gastric transit times were 0.63 h, 1.36 h, and 0.73 h for solution, capsules and tablets, respectively.<sup>8</sup> Other studies evaluating gastric emptying and food effects on oral drug absorption used Minipigs kept on straw bedding throughout the study,<sup>9,10</sup> a factor which likely influenced the findings in those studies.

The tolerability to many compounds is higher and more human-like in Göttingen Minipigs than in the dog. For instance, dogs show excessive vomiting in response to many compounds, including anticancer compounds,<sup>11</sup> and have a low tolerance to NSAIDs, that tend to result in gastrointestinal lesions.<sup>3</sup>

A recent review presented data demonstrating the suitability of the pig as a preclinical model to predict human oral bioavailability (with slightly better performance than the dog).<sup>5</sup> The overall similarity between humans and Minipigs regarding intestinal transporters and metabolizing enzymes in the intestine and liver,<sup>5,12</sup> helps to explain this finding.



*“Over the years, the large number of toxicological studies performed in Göttingen Minipigs at Scantox in Denmark have involved many different routes of administration. The oral dosing route is of the one most commonly used, and one that we have good experience with.”* Scantox, Denmark

**For more information, please contact:**

Ellegaard Göttingen Minipigs A/S  
Soroe Landevej 302  
4261 Dalmoose, Denmark  
Tel.: +45 5818 5818  
[ellegaard@minipigs.dk](mailto:ellegaard@minipigs.dk)





## References

1. Mößeler A, Vagt S, Beyerbach M, Kamphues J. Starch origin and thermal processing affect starch digestion in a minipig model of pancreatic exocrine insufficiency. *Gastroenterol Res Pract.* 2015;2015:872872
2. Bode G, Clausing P, Gervais F, et al. The utility of the minipig as an animal model in regulatory toxicology. *J Pharmacol Toxicol Methods.* 2010;62:196-220
3. Swindle MM, Makin A, Herron AJ, et al. Swine as models in biomedical research and toxicology testing. *Vet Pathol.* 2012;49:344-356
4. Suenderhauf C, Parrott N. A physiologically based pharmacokinetic model of the minipig: data compilation and model implementation. *Pharm Res.* 2013;30:1-15
5. Henze LJ, Koehl NJ, O'Shea JP, et al. The pig as a preclinical model for predicting oral bioavailability and in vivo performance of pharmaceutical oral dosage forms: a PEARRL review. *J Pharm Pharmacol.* 2018 Apr 10. doi: 10.1111/jphp.12912
6. Diehl KH, Hull R, Morton D, et al. A good practice guide to the administration of substances and removal of blood, including routes and volumes. *J Appl Toxicol.* 2001;21:15-23
7. Preuße C, Skaanild MT. Minipigs in absorption, distribution, metabolism, and excretion (ADME) studies. In McNulty P, Dayan AD, Ganderup NC, Hastings KL(Eds.), *The minipig in biomedical research* (pp. 143-158). 2012. Taylor & Francis, Boca Raton, FL
8. Suenderhauf C, Tuffin G, Lorentzen H, et al. Pharmacokinetics of paracetamol in Göttingen Minipigs: in vivo studies and modelling to elucidate physiological determinants of absorption. *Pharm Res.* 2014;31:2696-2707
9. Christiansen ML, Müllertz A, Garmer M, et al. Evaluation of the use of Göttingen Minipigs to predict food effects on the oral absorption of drugs in humans. *J Pharm Sci.* 2015;104:135-143
10. Henze LJ, Griffin BT, Christiansen M, et al. Exploring gastric emptying rate in minipigs: Effect of food type and pre-dosing of metoclopramide. *Eur J Pharm Sci.* 2018;118:183-190
11. Mahl A, Dincer Z, Heining P. The potential of minipigs in the development of anticancer therapeutics. *Toxicol Pathol.* 2016;44:391-397
12. Dalgaard L. Comparison of minipig, dog, monkey and human drug metabolism and disposition. *J Pharmacol Toxicol Methods.* 2015;74:80-92

---

### For more information, please contact:

Ellegaard Göttingen Minipigs A/S  
Soroe Landevej 302  
4261 Dalmoose, Denmark  
Tel.: +45 5818 5818  
ellegaard@minipigs.dk



[www.minipigs.dk](http://www.minipigs.dk)